
CONTENTS

Lizards and geckos	1
Primates	3

LIZARDS AND GECKOS

A study of home range, territoriality and activity level of *Zonosaurus laticaudatus* in the dry and semideciduous forest of Kirindy

Abstract

The study was conducted to determine home range, territoriality and the level of activity for different periods of the day. It also analyses the effect of size and sex on home range size. Twenty one individuals were marked (9 males, 11 females) and observed between 17 and 24 November 2002. The data showed no correlation between home range size with both individual body size and sex. The territories of different individuals showed little overlap suggesting territoriality. Overlap was more pronounced within the same sex than for both sexes. The data showed a variation in the level of activity and the type of activity performed for different times of day.

Samuel Ayesu, University of Sciences and Technology, Ghana

Razafindrakoto Germain Fanomezantsoa, University of Antananarivo, Madagascar

Toto Volahy Anselme, Durrell Wildlife Conservation Trust, Madagascar

2002

Survey of the nocturnal ground geckos of the genus *Paroedura* in the dry deciduous forest of Kirindy

Abstract

The nocturnal ground geckos of the endemic *Paroedura* genus are one of the lesser-known groups of reptiles of Madagascar. Not even the systematics has been studied cleared, new species are still steadily discovered. We tried to sort out the relationships within the *Paroedura* of Kirindy forest. Two species are easily to be distinguished *Paroedura bastardi* and *Paroedura picta*. But *P. picta* was found to occur in three phenotypes that differ in colour pattern and as we found also in other morphological characteristics. Some people believe these three types to be distinct species; our results can't clearly favour a pro or contra. We hope this study will contribute in understanding the morphological and ecological features of the *Paroedura* species of Kirindy. The structure within this genus is still unclear but Raxworthy & Nussbaum are working on the phylogeny of this genus at the moment.

Sven Selbert, Humboldt University Berlin, Germany

Jonah van Beijnen, Wageningen University, The Netherlands

2003

Rotten logs: their importance as a microhabitat for geckos in Kirindy Forest

Abstract

The main aim of this project was to assess the role of rotten logs in the Kirindy forest reserve as a microhabitat for geckos. Transects of 140 m by 20 m were sampled over six days for road, path and forest habitat types. Some physical parameters of the logs were also measured. 68 individuals belonging to seven genera and nine species were observed for 360 logs sampled. Results also showed that geckos use these logs irrespectively of the temperature, but primarily for protection, as they preferred being within vertical logs, allowing them to climb away from ground based predators.

Benjamin Isambert, Natural History Museum, UK

Rosemary Nanji, University of Yaounde I, Cameroon

Lalandy Razafy Sehen, University of Antananarivo, Madagascar

2006

Behavioural ecology of iguanid lizard *Oplurus cuvieri* in Kirindy forest: associations with sex and colour morph

Abstract

Kirindy dry deciduous forest is a Special Reserve in the central western region of Madagascar. *Oplurus cuvieri* an endemic species of iguanid (Oplurine) lizard found in the region was investigated in order to determine whether the natural colour variability bears an association with behaviour. The distribution of colour morphs (light, intermediate and dark) between sexes show that it is more likely to be a sex-linked trait rather than one which might affect behaviour or thermoregulation. Furthermore, our study found differences in behavioural activity patterns between males and females. In particular, males move greater distances than females as a facet of breeding season, while females hold smaller home ranges.

Harry Fox, Durham University, England

Jan Schnitzler, Royal Botanic Gardens, Kew, England

2007

Activity patterns of Ground Geckos (*Paroedura picta*) in Kirindy Forest

Abstract

Little is known about behaviour of nocturnal ground geckos. To augment existing knowledge on that topic, activity patterns of *Paroedura picta* were observed in Kirindy Forest, Madagascar, and mark-recapture methods were applied. In total 70 adult and 48 juvenile geckos were observed during the study. There was no difference between early and late night activities; between males and females and between adults and juveniles. The distance moved ranged between 0 and 25.8 m per night with a mean of 7.4 m. No preferences were observed either for soil type or for vegetation density.

Adam Cieplinski, University of Warsaw, Poland

Eleni Foui, University of Leeds, UK

Niklaus Peyer, University of Zurich, Switzerland

2008

PRIMATES

**The diversity of sifaka diet during a period of stress
in Kirindy forest, Madagascar****Abstract**

The dry deciduous forest of Kirindy on the west coast of Madagascar is home to the Verreaux's Sifaka (*Propithecus verreauxi verreauxi*). The seasonality of the climate affects the foraging behaviour of the Sifaka. We analysed the dietary aspects of the Sifaka in a six day period at the end of the dry season. Our results show no preference for specific plant parts (leaves vs. other). The diet consists largely of endemic species, but preference was not significant. The onset of precipitation at the end of the dry season did not cause a foraging switch in feeding time on lianas but did so for trees. Our study shows that the Sifaka has a species rich diet at the end of the dry season. The persistence of viable Sifaka populations is coupled to the high species richness of Kirindy Forest. Additionally, the forest holds numerous local endemic species. We therefore suggest a preservation, rather than a conservation strategy, for the future management of the reserve.

Anna Norman, Stockholm University, Sweden

Joeri S. Strijk, Université Paul Sabatier, France

2006
